arXiv:1209.2332v1 [hep-ph] 11 Sep 2012

Propagator poles and an emergent stable state below threshold: general discussion and the $E(38)$ state

Francesco Giacosa and Thomas Wolkanowski
Institut für Theoretische Physik, Johann Wolfgang Goethe - Universität
Max von Laue–Str. 1, D-60438 Frankfurt am Main, Germany

September 12, 2012

Abstract

In the framework of a simple quantum field theory describing the decay of a scalar state into two (pseudo)scalar ones we study the pole(s) motion(s) of its propagator: besides the expected pole on the second Riemann sheet, we find – for a large enough coupling constant – a second, additional pole on the first Riemann sheet below threshold, which corresponds to a stable state. We then perform a numerical study for a hadronic system in which a scalar particle couples to pions. We investigate under which conditions a stable state below the two-pion threshold can emerge. In particular, we study the case in which this stable state has a mass of 38 MeV, which corresponds to the recently claimed novel scalar state $E(38)$. Moreover, we also show that the resonance $f_0(500)$ and the stable state $E(38)$ could be two different manifestation of the same 'object'. Finally, we also estimate the order of magnitude of its coupling to photons.

In this Letter we study the propagator, its poles, and the spectral function of a scalar state S which interacts with a (pseudo)scalar field φ as described in the following Lagrangian [1, 2]:

$$\mathcal{L} = \frac{1}{2}(\partial_\mu S)^2 - \frac{1}{2}M_0^2 S^2 + \frac{1}{2}(\partial_\mu \varphi)^2 - \frac{1}{2}m^2\varphi^2 + gS\varphi^2 \ . \tag{1}$$

The propagator of the field S is dressed by loops of φ-particles. The contribution of one loop is evaluated using the standard Feynman rules,

$$\Sigma(p^2) = -i\int \frac{\mathrm{d}^4 q}{(2\pi)^4} \frac{\phi^2(q)}{\left[\left(\frac{p}{2}+q\right)^2 - m^2 + i\varepsilon\right]\left[\left(\frac{p}{2}-q\right)^2 - m^2 + i\varepsilon\right]} \ , \tag{2}$$

where the function $\phi(q)$ has been introduced to regularize the integral. (Formally, it can be included at the Lagrangian level by rendering it nonlocal, see Ref. [3] and refs. therein). Here we adopt the choice $\phi(q) = \tilde{\phi}(\mathbf{q}^2) = \left(1 + \mathbf{q}^2/\Lambda^2\right)^{-1}$ [2], where Λ sets the energy scale at which the loop contribution gets suppressed. It should be stressed that the precise form of the cutoff function affects the results only marginally, as long as it falls off sufficiently fast to guarantee convergence.

Upon resummation of the one-loop contributions, the propagator of S takes the form

$$G_S(p^2) = \left[p^2 - M_0^2 + (\sqrt{2}g)^2\Sigma(p^2) + i\varepsilon\right]^{-1} \ . \tag{3}$$

The spectral function of the state S is obtained as the imaginary part of the propagator as [2, 4, 5]

$$d_S(x = \sqrt{p^2}) = \frac{2x}{\pi}\left|\lim_{\varepsilon\to 0^+} \mathrm{Im}[G_S(x^2)]\right| \ . \tag{4}$$

As a consequence of the Källen–Lehmann representation, the normalization $\int_0^\infty d_S(x)\mathrm{d}x = 1$ holds.

We first describe the general qualitative features of our results, which are independent on numerical details. We assume that the (bare) mass M_0 lies well above the threshold $2m$, thus implying that the decay into $\varphi\varphi$ is kinematically allowed. As soon as the coupling constant g does not vanish, the propagator (3) is a multivalued analytic function for the complex variable $z = \sqrt{p^2}$. The resonance corresponding to the field S is related to a pole on the (unphysical) second Riemann sheet. We denote this pole as

$$z_{\mathrm{II}} = M_{pole} - i\Gamma_{pole}/2 \ , \tag{5}$$

where M_{pole} is usually referred to as the pole mass of the resonance S.

Clearly, for $g \to 0$ one has $M_{pole} \to M_0$ and $\Gamma_{pole} \to 0$. When varying the coupling constant g a trajectory in the complex plane is generated, see the dashed line in Fig. 1 for a numerical example and the discussion below. Note, for small values of g the decay width Γ_{pole} obtained from Eq. (5) is reproduced very well by the tree-level result $\Gamma_{S\varphi\varphi}(M_{pole})$, where the tree-level decay function $\Gamma_{S\varphi\varphi}(x)$ reads $\Gamma_{S\varphi\varphi}(x) = \frac{\sqrt{x^2/4-m^2}}{8\pi x^2}(\sqrt{2}g)^2\tilde{\phi}^2(x^2/4-m^2)$. In this regime the spectral function $d_S(x)$ is well approximated by a Breit–Wigner function, see Fig. 2 (solid curve) for a typical Breit–Wigner shaped resonance. However, for increasing coupling (even sizable) distortions arise [2]. Note, departures from Breit–Wigner also imply a non-exponential decay when the time evolution of the unstable state is studied [6]. This is the quantum field theoretical counterpart of a general quantum phenomenon, see for example Refs. [8, 9].

Besides the pole (5), other poles may exist on the II-Riemann sheet, which however have practically no influence on the determination of the properties of the resonance S. Interestingly, for moderate values of the coupling constant g there is a pole on the real axis below threshold. This pole, although not important for the phenomenology of the resonance, moves for increasing g along the real axis towards the threshold value $2m$. Then, for the coupling constant g exceeding a critical value g_*, this pole disappears from the II-Riemann sheet, but appears on the real axis below threshold on the I-Riemann sheet. We denote this newly emerged pole on the first sheet as

$$z_{\mathrm{I}} = \tilde{M} - i\varepsilon \text{ with } \tilde{M} < 2m \text{ (for } g \geq g_* \text{ only).} \tag{6}$$

The pole z_{I} changes considerably the physical properties of the system: namely, it corresponds to a stable state below threshold. Thus, for $g \geq g_*$ there are two relevant poles on the Riemann manifold: the standard resonance pole on the second sheet of Eq. (5) and the additional ‘stable state’ pole from Eq. (6) on the first sheet. The spectral function in the presence of the additional pole below threshold takes the form

$$d_S(x) = Z\delta(x - \tilde{M}) + d_{a.t.}(x) \ , \tag{7}$$

where $d_{a.t.}(x)$ (a.t. stands for above threshold) vanishes for $x < 2m$. The quantity Z is nonzero for $g \geq g_*$ and its explicit form is given by

$$Z = \left[1 + (\sqrt{2}g)^2 \left(\frac{\partial\Sigma(x^2)}{\partial x^2}\right)_{x^2=\tilde{M}^2}\right]^{-1} \ . \tag{8}$$

The full spectral function $d_S(x)$ is still normalized to 1, implying that $Z + \int_{2m}^{\infty} d_{a.t.}(x)\mathrm{d}x = 1$.

The emergence of additional poles was described in the literature (usually connected to light scalar mesons), e.g. Refs. [10, 11, 12, 13]. As described in detail in Ref. [13], besides a massive ‘seed’ pole, a second pole on the second Riemann sheet with a smaller real part exists. The movement of this pole in the complex plane is peculiar: for small values of (their) coupling constant, its imaginary part is very large. Then, by increasing the coupling, both the imaginary and the real part of this additional pole become smaller. Eventually, for a coupling large enough, a stable state below threshold is obtained. It should be stressed that the pole we obtain out of our Lagrangian (1) has very different properties: namely, as described above, it is situated on the real axis of the second sheet for small and intermediate values of the coupling, then vanishes for a coupling large enough by slipping into the branch cut and appears on the real axis of the first sheet.



Figure 1: The trajectory (dashed line) of the second-sheet pole z_{II} (Eq. (5)) obtained by varying the coupling constant g is shown for the numerical values in Eq. (9). For some examples (white markers along the dashed line) the value of g is reported. When existent, we have also shown – using the same form of the markers – the first-sheet pole z_{I} (Eq. (6)) lying on the real axis below threshold. For instance, in the case $g = g_E$ the pole z_{I} corresponds to a stable state with the mass $\tilde{M} = 38$ MeV.

We now turn to some numerical calculations in order to show some explicit examples of the described properties. For illustrative purposes we use the following numerical values, which are typical in hadronic theories [14]:

$$M_0 = 1 \text{ GeV} \ , \ \Lambda = 1.5 \text{ GeV} \ , \ m = m_{\pi^0} = 0.135 \text{ GeV} \ . \tag{9}$$

Thus, the threshold $2m$ corresponds to the minimal energy threshold of QCD, namely that of two pions. Note, the choice of $M_0 = 1$ GeV in Eq. (9) is not linked to a resonance listed in the Particle Data Group (PDG) [15]. It corresponds to a hypothetical resonance with a mass of about 1 GeV which couples strongly to pions. This resonance is supposed to be a hadronic state made of quarks and gluons (eventually a broad tetraquark state), but in the present framework it serves just as a tool to test our approach.

When varying the value of the coupling constant g from small to large values, the position of the resonance pole z_{II} defines a trajectory on the II-Riemann sheet, which is depicted in Fig. 1 (dashed line). Moreover, for $g \geq g_* = 3.739$ GeV the first-sheet (alias bound-state) pole $z_{\mathrm{I}} = \tilde{M} - i\varepsilon$ is also present and its position for some values of g is shown.

In particular, we have shown in Fig. 1 also the case $g = g_E = 4.877 \text{ GeV} > g_*$ for which the mass of the stable state on the first sheet reads $z_{\mathrm{I}} = \tilde{M} - i\varepsilon = (38 - i\varepsilon)$ MeV. Quite interestingly, in Ref. [16] E. van Beveren and G. Rupp discuss the possible existence of a new boson with a mass of 38 MeV, called $E(38)$, by a reanalysis of BABAR data [17] for the process $e^+e^- \to \Upsilon(1, 2^3S_1)\pi^+\pi^- \to l^+l^-\pi^+\pi^-$ where Υ stands for a bottonium state and l for a lepton. Namely, they argue that a light bosonic state with a mass of 38 MeV is emitted from the decay of the bottonium, resulting in a missing mass for the process described above. This hypothetical boson with a mass of 38 MeV then further decays into photons. In Ref. [18] the same authors discuss the data of the COMPASS experiment [19], which allegedly add evidence in favour of the existence of $E(38)$ and confirms the decay $E(38) \to \gamma\gamma$. This results has been disputed in a comment by the COMPASS collaboration [20]; later on, E. van Beveren and G. Rupp replied to the comment [21]. Recently, experimental results from the JINR Nuclotron



Figure 2: The spectral function of the state S for two different values of the coupling constant is shown for the numerical values in Eq. (9). The solid (blue) curve corresponds to $g = 1.5$ GeV: the second-sheet pole is $z_{\text{II}} = (0.985 - i0.038)$ GeV. A typical Breit–Wigner form is visible. The dashed (red) curve corresponds to $g = g_E$, for which the second-sheet pole $z_{\text{II}} = (1.096 - i0.472)$ GeV lies deep in the lower half plane. Being $g_E > g_*$, there is also the firs-sheet pole $z_{\text{I}} = (0.038\ - i\varepsilon)$ GeV. The spectral function shows a stable δ-peak for 0.038 GeV and a broad enhancement above threshold.

from DUBNA claiming the verification of the existence of the $E(38)$ state has been reported in Ref. [22], in which (utterly different) nuclear reactions involving two photons in the final state have been studied.

It should be stressed that the issue about the existence of $E(38)$ is under discussion and confirmation is needed; nevertheless, due to the fact that within our simple quantum field theoretical approach we can naturally accommodate a pole below threshold, it is amusing and interesting to investigate this numerical example in more details. We thus show in Fig. 2 the spectral function of the state S for two values of g. For the small value $g = 1.5$ GeV there is no pole on the first Riemann sheet and the spectral function has a typical 70 MeV broad Breit–Wigner form centered at about $M_0 = 1$ GeV. For the case $g = g_E$ the pole z_{I} on the first sheet is present. The spectral function has the form of Eq. (7) with $Z = 0.28$. Below threshold there is a δ-type peak for $\tilde{M} = 38$ MeV, above threshold there is a very broad and flat enhancement centered at about 1 GeV. It is clear that such a structure would be very hard to detect: its influence on $\varphi\varphi$ (alias pion-pion) scattering is expected to be very small, thus also explaining why this hypothetical broad state could not be observed experimentally. Indeed, the coupling of S to pions is large but not unrealistic: $g_\pi = g_E/\sqrt{3} = 2.816$ GeV, where $\sqrt{3}$ takes into account isospin symmetry, not included in our Lagrangian. A possibility could be a broad tetraquark state, for which the coupling constant is of the same order [23, 24].

A quite interesting and remarkable result follows when testing other values for M_0. Namely, one of the most famous mesonic resonance is the lightest scalar meson $f_0(500)$. In the new version of the PDG the pole of this resonance is estimated as (0.40-0.55) $- i$(0.20-0.35) GeV [15]. It is then natural to ask if we could assign the $f_0(500)$ resonance to our state S and at the same time describe the stable state $E(38)$. Indeed, by simply changing the mass to $M_0 = 0.45$ GeV we obtain the first-sheet pole $z_{\text{I}} = (0.038 - i\varepsilon)$ GeV for $g = g_E = 2.189$ GeV. The second-sheet pole reads $z_{\text{II}} = (0.427 - i0.371)$ GeV, which is *remarkably* close to the range given above. We summarize the results of this study in Fig. 3, in which both the z_{II} pole trajectory and the spectral function for $g = g_E$ are shown. It is therefore conceptually possible to link the states $f_0(500)$ and $E(38)$. It is worth to mention that there are indeed phenomenological models in which the state $f_0(500)$ is interpreted as a tetraquark state and



Figure 3: Left: z_{II} pole trajectory for $M_0 = 0.45$ GeV. The critical value is here given by $g_* = 1.399$ GeV. For $g = g_E = 2.189$ GeV the first-sheet pole $z_{\mathrm{I}} = (0.038 - i\varepsilon)$ GeV is obtained. In the shadowed rectangular the ranges of the PDG estimate for the $f_0(500)$ pole is reported: it is visible that the z_{II} pole (white square) lies very close to it. Right: spectral function $d_S(x)$ for the same numerical values. Here $Z = 0.68$, showing that the pole below threshold gives the dominant contribution to the spectral function.

which reduce to a Lagrangian of the type (1) when only the pion-pion channel is studied [23]. Surely, more advanced investigations in this direction should go beyond our simple model, test different cutoff functions and take into account other interaction forms, such as the inclusion of derivatives [25]. Most important, one should also include chiral symmetry. Nevertheless the possibility to describe $f_0(500)$ and $E(38)$ in an unified framework is intriguing.

As a last subject of this work we describe the coupling to photons. To this end we add to the Lagrangian the following interaction term:

$$\mathcal{L}_{\gamma\gamma} = c_{\gamma\gamma} S F_{\mu\nu} F^{\mu\nu} \tag{10}$$

where $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$ and A^μ is the photon field. The tree-level decay width (as function of the running mass x) reads $\Gamma_{S\to\gamma\gamma}(x) = c_{\gamma\gamma}^2 x^3/4\pi$. Now, $c_{\gamma\gamma}$ is surely not zero because S, being made of quarks, can couple also to photons. However, its precise value is unknown. Usually, the typical order of magnitude for $\gamma\gamma$ decays of mesonic states amounts to some keV.

For $g \geq g_*$ there is, as described above, a (hadronic) stable state below threshold. This state can however decay into photons. In order to evaluate the $\gamma\gamma$ decay width the redefinition $S \to \sqrt{Z} S$ is necessary to guarantee the correct residuum of the propagator. Referring for example to the case $g = g_E$ with $z_{\mathrm{I}} = (0.038 - i\varepsilon)$ GeV we find for the numerical values of Eq. (9):

$$\Gamma_{E(38)\to\gamma\gamma} = Z c_{\gamma\gamma}^2 \frac{\tilde{M}^3}{4} = Z \left(\frac{\tilde{M}}{M_0}\right)^3 \Gamma_{S\to\gamma\gamma}(M_0 = 1 \text{ GeV}) = 1.55 \cdot 10^{-5}\ \Gamma_{S\to\gamma\gamma}(M_0 = 1 \text{ GeV}) \ . \tag{11}$$

Assuming that $\Gamma_{S\to\gamma\gamma}(M_0 = 1 \text{ GeV})$ is of the order of 10 keV, we obtain that

$$\Gamma_{E(38)\to\gamma\gamma} \simeq 10^{-4} \text{ keV} \ , \tag{12}$$

which is a very small value. If, instead, we study the case of $M_0 = 0.45$ GeV and $g = g_E = 2.189$ GeV we obtain $\Gamma_{E(38)\to\gamma\gamma} = 4.1 \cdot 10^{-4} \Gamma_{f_0(500)\to\gamma\gamma}$. Using the value $\Gamma_{f_0(500)\to\gamma\gamma} \simeq 1$ keV (see the list of results in [15]) one gets $\Gamma_{E(38)\to\gamma\gamma} \simeq 4.1 \cdot 10^{-4}$ keV, which is similar to the previous estimate.

In conclusions, we have presented a simple quantum field theoretical model involving scalar particles and studied the poles in the first and second Riemann sheet. We have shown that for a coupling constant large enough one obtains, in addition to the standard resonance pole on the second sheet, also a pole below threshold on the first sheet. The latter corresponds then to an emergent stable state. We have studied this system using the numerical values typical for a hadronic system and have shown that the appearance of the additional stable state is realistic (Fig. 1 and 2). In particular, we have studied the case in which a hadronic stable resonance with a mass of 38 MeV is realized: this is interesting in view of recent claims of the existence of such a state, called $E(38)$. Indeed, it is also possible that the $E(38)$ and the resonance $f_0(500)$ (alias σ) are two manifestations of the same object (Fig. 3). In the end, we have explored the decay of $E(38)$ into two photons. The present theoretical description of the putative $E(38)$ state is very different from the one put forward in Refs. [16, 18, 26]. Thus, our analysis offers a new point of view which may help in the process of falsification of what this state, if existent, is or is not.

Acknowledgments: The authors thank George Rupp, Giuseppe Pagliara and Dirk H. Rischke for valuable discussions. In particular, George Rupp is also acknowledged for useful comments about the $E(38)$ state. F.G. thanks the Foundation Polytechnical Society Frankfurt am Main for support through an Educator fellowship.

References

[1] M. J. G. Veltman, Physica **29** (1963) 186.

[2] F. Giacosa and G. Pagliara, Phys. Rev. C **76** (2007) 065204 [arXiv:0707.3594 [hep-ph]].

[3] J. Terning, Phys. Rev. D **44** (1991) 887. G. V. Efimov and M. A. Ivanov, "The Quark confinement model of hadrons," *Bristol, UK: IOP (1993) 177 p.* Y. V. Burdanov, G. V. Efimov, S. N. Nedelko and S. A. Solunin, Phys. Rev. D **54** (1996) 4483 [arXiv:hep-ph/9601344]. A. Faessler, T. Gutsche, M. A. Ivanov, V. E. Lyubovitskij and P. Wang, Phys. Rev. D **68** (2003) 014011 [arXiv:hep-ph/0304031]. F. Giacosa, T. Gutsche and A. Faessler, Phys. Rev. C **71** (2005) 025202 [arXiv:hep-ph/0408085].

[4] P. T. Matthews and A. Salam, Phys. Rev. **112** (1958) 283. P. T. Matthews and A. Salam, Phys. Rev. **115** (1959) 1079.

[5] N. N. Achasov and A. V. Kiselev, Phys. Rev. D **70** (2004) 111901 [arXiv:hep-ph/0405128].

[6] F. Giacosa and G. Pagliara, Mod. Phys. Lett. **A26** (2011) 2247-2259. [arXiv:1005.4817 [hep-ph]]. G. Pagliara, F. Giacosa, Acta Phys. Polon. Supp. **4** (2011). [arXiv:1108.2782 [hep-ph]].

[7] F. Giacosa, Found. Phys. **42** (2012) 1262 [arXiv:1110.5923 [nucl-th]].

[8] L. Fonda, G. C. Ghirardi and A. Rimini, Rept. Prog. Phys. **41** (1978) 587.

[9] P. Facchi, H. Nakazato, S. Pascazio Phys. Rev. Lett. **86** (2001) 2699-2703.

[10] E. van Beveren, T. A. Rijken, K. Metzger, C. Dullemond, G. Rupp and J. E. Ribeiro, Z. Phys. C **30** (1986) 615 [arXiv:0710.4067 [hep-ph]].

[11] N. A. Tornqvist, Z. Phys. C **68** (1995) 647 [hep-ph/9504372].

[12] M. Boglione and M. R. Pennington, Phys. Rev. D **65** (2002) 114010 [hep-ph/0203149].

[13] E. van Beveren, D. V. Bugg, F. Kleefeld and G. Rupp, Phys. Lett. B **641** (2006) 265 [hep-ph/0606022].

[14] C. Amsler and N. A. Tornqvist, Phys. Rept. **389**, 61 (2004). E. Klempt and A. Zaitsev, Phys. Rept. **454** (2007) 1

[15] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86** (2012) 010001.

[16] E. van Beveren and G. Rupp, arXiv:1102.1863 [hep-ph].

[17] B. Aubert *et al.* [BABAR Collaboration], of Upsilon(4S) —¿ eta Upsilon(1S) decay," Phys. Rev. D **78** (2008) 112002 [arXiv:0807.2014 [hep-ex]].

[18] E. van Beveren and G. Rupp, arXiv:1202.1739 [hep-ph].

[19] J. Bernhard *et al.* [COMPASS Collaboration], arXiv:1109.0272 [hep-ex]. T. Schluter [COMPASS Collaboration], eConf C **110613** (2011) 83 [arXiv:1108.6191 [hep-ex]].

[20] [COMPASS Collaboration], arXiv:1204.2349 [hep-ex].

[21] E. van Beveren and G. Rupp, arXiv:1204.3287 [hep-ph].

[22] K. .U. Abraamyan, A. B. Anisimov, M. I. Baznat, K. K. Gudima, M. A. Nazarenko, S. G. Reznikov and A. S. Sorin, arXiv:1208.3829 [hep-ex].

[23] F. Giacosa, Phys. Rev. D **74** (2006) 014028 [hep-ph/0605191].

[24] L. Maiani, F. Piccinini, A. D. Polosa and V. Riquer, Phys. Rev. Lett. **93** (2004) 212002 [arXiv:hep-ph/0407017]. G. 't Hooft, G. Isidori, L. Maiani, A. D. Polosa and V. Riquer, Phys. Lett. B **662** (2008) 424 [arXiv:0801.2288 [hep-ph]].

[25] F. Giacosa and G. Pagliara, Nucl. Phys. A **812** (2008) 125 [arXiv:0804.1572 [hep-ph]].

[26] C. Dullemond, T. A. Rijken and E. van Beveren, Nuovo Cim. A **80** (1984) 401.